

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

INNS011/KAM

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/297981INTERNATIONAL APPLICATION NO.
PCT/EP98/05518INTERNATIONAL FILING DATE
31 August 1998PRIORITY DATE CLAIMED
29 August 1997

TITLE OF INVENTION

Methylated, SmD Homologous Peptides, Reactive with the Antibodies From Sera of Living Beings Affected With Systemic Lupus Erythematosus

APPLICANT(S) FOR DO/EO/US

MEHEUS, Lydie; LÜHRMANN, Reinhard, Georg; UNION, Ann; RAYMACKERS, Joseph

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 30(1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11 to 16 below concern document(s) or information included:

11. ☒ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☐ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☒ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

Postcard, Fee Calculation Sheet (in duplicate); check for \$1,608.00, Petition for Revival and check for \$1210.00.

FEE VALUE ACCOUNT/ABILITY	
DEPOSIT	ACCOUNT NO.
01	2508
FEE CODE	VALUE FURNISHED
910	970
970	840

CERTIFICATE OF EXPRESS MAILING

NUMBER **EL291389138US**DATE OF DEPOSIT **May 10, 1999**

This paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Assistant Commissioner for Patents, Washington, DC 20231

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)	INTERNATIONAL APPLICATION NO. PCT/EP98/05518	ATTORNEY'S DOCKET NUMBER INNS011/KAM
---	---	---

17. ☒ The following fees are submitted:

Basic National Fee (37 CFR 1.492(a)(1)-(5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO.....\$970.00

International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO.....\$840.00

International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....\$760.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
but all claims did not satisfy provisions of PCT Article 33(1)-(4).....\$670.00

International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4).....\$ 96.00

ENTER APPROPRIATE BASIC FEE AMOUNT = \$970.00

Surcharge of \$130.00 for furnishing the oath or declaration later than ☒ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(e)). **\$130.00**

Claims	Number Filed	Number Extra	Rate
Total Claims	19 - 20 =	0	x \$ 18.00
Independent Claims	9 - 3 =	6	x \$ 78.00
Multiple dependent claim(s) (if applicable)			+ \$260.00

TOTAL OF ABOVE CALCULATIONS = \$ 468.00

Reduction by ½ for filing by small entity, if applicable. Verified Small Entity statement must
also be filed. (NOTE: 37 CFR 1.9, 1.27, 1.28) **\$ 1568.00**

SUBTOTAL = \$.00

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)). **\$.00**

TOTAL NATIONAL FEE = \$.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property) **\$ 40.00**

TOTAL FEES ENCLOSED = \$1,608.00

	Amount to be refunded:	\$.00
	charged	\$.00

CALCULATIONS
PTO USE ONLY

a. ☒ A check in the amount of \$ 1,608.00 to cover the above fees is enclosed. Applicants request a Notice of Missing Parts to satisfy the filing of the Declaration and Preliminary Amendment.


b. ☐ Please charge my Deposit Account No. 01-2508/ in the amount of \$ to cover the above fees. A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 01-2508/INNS011/KAM. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

Patricia A. Kammerer
ARNOLD WHITE & DURKEE
P.O. Box 4433
Houston, TX 77057-2198
(713) 787-1400


SIGNATURE

Patricia A. Kammerer
NAME

29,775
REGISTRATION NUMBER

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

LYDIE MEHEUS
REINHARD GEORG LÜHRMANN
ANN UNION
JOSEPH RAYMACKERS

Group Art Unit:

Examiner:

Atty. Dkt. No.: INNS011/KAM

Serial No.:

Filed:

For: METHYLATED, SmD HOMOLOGOUS
PEPTIDES, REACTIVE WITH THE
ANTIBODIES FROM SERA OF LIVING
BEINGS AFFECTED WITH SYSTEMIC
LUPUS ERYTHEMATOSUS

Int'l Application No. PCT/EP98/05518

Int'l Filing Date: August 31, 1998

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

CERTIFICATE OF EXPRESS MAILING	
NUMBER	EL291389138US
DATE OF DEPOSIT:	<i>May 10, 1999</i>
This paper or fee is being deposited with the United States Postal Service "EXPRESS MAIL POST OFFICE TO ADDRESSEE" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to Assistant Commissioner for Patents, Washington, DC 20231.	

Please amend this application as follows:

IN THE SPECIFICATION:

At page 3 line 30, insert the heading

--SUMMARY OF THE INVENTION--;

At page 7, line 8, insert the heading

--DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS--;

At page 31, line 6 please delete the heading "Figure Legends";

At page 31, lines 8-27 delete the text and insert it instead at page 7, line 7, and preceded by the caption --BRIEF DESCRIPTION OF THE FIGURES--.

IN THE CLAIMS:

Please cancel without prejudice claims 16-18 and amend claims 1, 3-15 and 19-22 as follows:

1. (Amended) Peptide containing less than 50 amino acids, comprising at least one dimer of the type XG, wherein X stands for a N^G-mono- or N^G-N^G-dimethylated arginine or N^G-N^G-dimethylated arginine, that is able to react with antibodies and with said methylation being crucial for the reaction between said peptide and said antibodies and wherein said antibodies are present in sera from patients with:

systemic lupus erythematosus, or

infectious, recurrent or chronic mononucleosis or infection, or

certain cancers which are related to infection with Epstein-Barr virus[, such as Burkitt's lymphoma or nasopharyngeal carcinoma].

3. (Amended) Peptide [and/or chemical structure comprising any of the peptides according to claims 1 or 2,] of claim 1 fused to a linker molecule.

4. (Amended) Circularized peptide that comprises at least one of the peptides according to claim 1 [any of the claims 1 to 3].

5. (Amended) Peptide comprising [and/or consisting of] tandem repeats of at least two of any of the peptides of claim 1 [of claims 1 to 4].

6. (Amended) Branched peptide that comprises at least one of the peptides according to claim 1 [any of the claims 1 to 5].

7. (Amended) Method for producing a peptide according to claim 1 [any of claims 1 to 6], by classical chemical synthesis, wherein methylated arginines are substituted for unmethylated arginine residues during the chemical synthesis.

8. (Amended) Method for producing a peptide according to claim 1 [any of claims 1 to 6], wherein the primary amino acid sequence is produced by classical chemical synthesis, and wherein the arginine residues that precede glycine residues are subsequently methylated by contacting said peptide with a protein arginine methyltransferase.

9. (Amended) Method for producing a peptide of [any of claims 1 to 6] claim 1 comprising the following steps:

transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements such that said peptide or a protein comprising said peptide is expressed and/or secreted,

culturing said transformed cellular host under conditions allowing expression of said protein or peptide and optionally allowing a partial or optimal methylation of the arginines present in said peptide,

harvesting said peptide.

10. (Amended) Method of claim 9 [for producing a peptide of any of claims 1 to 6] comprising the further step of [following steps:

transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements, such that said peptide or a protein comprising said peptide is expressed and/or secreted,

culturing said transformed cellular host under conditions allowing expression of said protein or said peptide,

harvesting said protein or said peptide,]

methylation arginine residues of said [protein or said] harvested peptide by contacting with a protein arginine methyltransferase.

11. (Amended) Method according to claim 9 [any of claims 9 or 10], wherein said host cell is a bacterial host or yeast or any other eukaryotic host cell which is preferably transformed with a recombinant baculovirus.

12. (Amended) An antibody raised upon immunization with a peptide according to claim 1 [any of the claims 1 to 6,] with said antibody being specifically reactive with the methylated forms of said peptide, and with said antibody being preferably a monoclonal antibody.

13. (Amended) Anti-idiotypic antibody raised upon immunization with an antibody according to claim 12, with said anti-idiotypic antibody being specifically reactive with the antibody of claim 12[, thereby mimicking the methylated form of a peptide according to any of claims 1 to 6, and with said antibody being preferably a monoclonal antibody].

14. (Amended) An immunotoxin molecule comprising and/or consisting of cell recognition molecule being a peptide of claim 1 [any of claims 1 to 6], or an antibody thereof [according to any of the claims 12 or 13], covalently bound to a toxin molecule or active fragment thereof.

15. (Amended) A medicament comprising a peptide according to claim 1, [any of the claims 1 to 6] or an antibody to said peptide, [according to any of claims 12 or 13] or an immunotoxin molecule comprising a toxin molecule covalently bound to said peptide or said antibody [according to claim 14 or a composition thereof for use as a medicament].

19. A diagnostic kit for use in detecting auto-immune diseases such as:

systemic lupus erythematosus,

discoid lupus erythematosus,

scleroderma,

dermatomyositis,

rheumatoid arthritis,

Sjögren's syndrome,

or for detecting diseases in which Epstein-Barr can be implicated such as:

Burkitt's lymphoma,

nasopharyngeal carcinoma,

Hodgkin's disease,

infectious, recurrent or chronic mononucleosis,

said kit comprising at least one peptide according to claim 1 [any of claims 1 to 6], or an antibody thereof [according to claims 12 or 13,] with said peptide or antibody [being possibly] optionally bound to a solid support.

20. A diagnostic kit according to claim 19, said kit comprising a range of peptides according to claim 1 [any of claims 1 to 6] or of antibodies thereof [according to claims 12 or 13, possibly] optionally in combination with native methylated SmD1 or SmD3 and recombinant unmethylated SmD1 or SmD3, wherein said peptides are attached to specific locations on a solid substrate.

21. A diagnostic kit according to claim 19 [or 20,] wherein said solid support is a membrane strip and said [poly]peptides are coupled to the membrane in the form of parallel lines.

[natural SmD (1,2 or 3) or in vitro dimethylated SmD (1, 2 or 3)

unmethylated SmD expressed in E.coli (1, 2 or 3)

peptide of any of claims 1 to 6]

22. A diagnostic kit according to any of claims 19 [to 21] wherein certain peptides are not attached to a solid support but are provided in the binding solution to be used as competitors and/or to block other antibodies that are present in sera from patients with autoimmune disease other than SLE, thereby decreasing or eliminating possible cross-reaction and/or aspecific binding.

Please add new claim 23:

--23. The peptide according to claim 1 wherein X stands for N^G-mono- or N^G-N^G-dimethylated arginine.--

REMARKS

The specification has been amended to conform to United States preferred arrangement in framing the specification. The claims have been amended to delete recitation of multiple dependency and clarify the claimed subject matter. Claim 1 has been amended to include coverage of N^G-N^G-dimethylated arginine. Support is found the amendment is found at page 10 lines 27-28. New claim 23 is added and finds support in original claim1. No new matter has been entered. Claims 1-15 and 19-23 are now pending.

Conclusion

In view of the foregoing amendments, applicants respectfully submit the claims are in proper form and condition for allowance. Applicants request that the claims be allowed and the application advanced to issue.

The Examiner is invited to contact the undersigned attorney at (713) 787-1438 with any questions, comments or suggestions relating to the referenced patent application.

Respectfully submitted,



Patricia A. Kammerer
Reg. No. 29,775
Attorney for Assignee
INNOGENETICS N.V.

ARNOLD, WHITE & DURKEE
P.O. Box 4433
Houston, Texas 77210-4433
(713) 787-1400

Date: May 10, 1999

Methylated, SmD homologous peptides, reactive with the antibodies
from sera of living beings affected with systemic lupus erythematosus.

5 The present invention relates to a method of producing certain peptides
containing methylated arginines that are followed by a glycine residue and that
constitute immunogenic determinants of antibodies present in sera from patients
with systemic lupus erythematosus, or Epstein-Barr virus and wherein the
methylation is a prerequisite for reacting with said antibodies. The invention also
10 relates to the use of said peptides for diagnosis and treatment of systemic lupus
erythematosus and related diseases, diseases in which Epstein-Barr virus has been
implicated.

Background of the invention

15 Systemic lupus erythematosus is an autoimmune disease, in which the
patient develops antibodies that react with many tissues of his own body.
Dominant antibodies are directed against components of the cell nucleus, with
epitopes that may be found in DNA, and in proteins that constitute small
ribonucleoprotein particles (snRNPs).

20 The first laboratory test ever devised for this disease was the LE (lupus
erythematosus) cell test. This test has to be repeated many times, before it results
in a positive reaction in about 90 % of the people with systemic lupus
erythematosus. Also, the LE cell test is not specific for lupus, and can be positive
in up to 20 % of the people with rheumatoid arthritis, in some patients with other
25 rheumatic conditions like Sjögren's syndrome or scleroderma, in patients with liver
disease, and in persons taking drugs such as hydralazine and procainamide. The
ANA test, which detects antibodies against nuclear antigens, is more specific for
lupus than the LE test, and is positive in many patients that suffer from systemic
lupus erythematosus. As with the LE test, a positive ANA is not diagnostic for
30 lupus since the test may also be positive in people with scleroderma,
dermatomyositis, rheumatoid arthritis, Sjögren's syndrome, in patients treated

with certain drugs, or in patients suffering from infectious mononucleosis, liver disease, malaria etc.

For these reasons and because the summed tests are expensive, new tests have been developed which are very helpful in the diagnosis of SLE. These include the anti-DNA antibody test, the anti-Sm antibody test, the anti-RNP antibody test, the anti-Ro antibody test, and tests which measure serum complement levels. Often, correct diagnosis will depend on the interpretation of many separate tests and symptoms.

The Sm antigen is a complex macromolecular structure consisting of 8 proteins (B, B', D1, D2, D3, E, F, G) associated with the U series of small RNA molecules. SmBB' and SmD are considered as the major antigenic components of the complex (for review see S.O.Hoch, 1989). However, SmBB' shows cross reactivity with the anti-RNP antibodies, consequently SmD is regarded as the most specific autoantigen for Sm (W.J. van Venrooij *et al.*, 1991).

The SmD cDNA has been isolated from a human B-lymphocyte library with synthetic oligonucleotide probes, designed on the basis of the N-terminal sequence of SmD (Rokeach *et al.*, 1988). Subsequently, it was shown that the *in vitro* transcription product could be immunoprecipitated by anti-Sm IgG. The D protein has since been characterized either as a doublet designated D and D' (Andersen *et al.*, 1990) or as three polypeptides designated D1 (16 kDa), D2 (16.5 kDa) and D3 (18 kDa) (Lehmeier *et al.*, 1990), D1 being identical with the SmD cloned by Rokeach *et al.* (1988). The sequence of D2 and D3 is substantially different from D1.

Over the years, several research groups have reported on the use of recombinant SmD and of SmD derived peptides and have published conflicting data. Rokeach *et al.* (1992a) expressed SmD1 in *E. coli* and in *S. cerevisiae*, but in contrast to the reactivity of natural SmD from HeLa cells, most of the patient anti-SmD sera bound recombinant SmD1 at a level not significantly higher than normal human sera. Nevertheless, the same group (Rokeach *et al.*, 1992b) has performed epitope mapping based on multiple fusions between the TrpE gene and fragments of the SmD coding sequence, expressed in *E. coli*. Two patterns of anti-

Sm reactivity emerged: discontinuous epitopes were found scattered over the full-length antigen, and a dominant epitope was located at the C-terminus, from amino acid 87 to 119 (Rokeach et al., 1992b). Using synthetic peptides, Barakat et al. (1990) showed that the N-terminus (peptide 1-20) and peptide 44-67 could be used as a valuable probe for SLE diagnosis although their results did not match the anti-SmD reactivity obtained by the traditional assay (patent EP-B-0491014). Using a similar strategy, Sabbatini et al. (1993a) have identified a dominant epitope in the C-terminal region of SmD1 (aa95-aa119) confirming the results of Rokeach et al. (1992b), but opposing the results obtained by Barakat et al. (1990). The most recent work on epitope mapping of SmD1 by means of synthetic peptides (James et al. 1994) showed that 8 of 9 SmD positive sera (precipitin positive) are reactive with the sequence spanning octapeptides 92-112. An additional epitope, clearly reactive with 7 of 9 SmD positive sera was located in the region of amino acid 82-90. Finally, a SmD-like epitope was recently identified by Rivkin et al. (1994) and consists of a (Gly-Arg)₉ dipeptide repeat (homology with the C-terminus). In contrast to the SLE specificity of anti-Sm antibodies, the defined epitope is also recognized by patients with other autoimmune diseases (rheumatoid arthritis, scleroderma, Sjögren's syndrome). The β -galactosidase fusion protein in *E.coli* of the above mentioned epitope was reactive with 35% of the SLE sera, but only 6 out of these 32 positive sera were reactive with the native SmD protein indicating that the fusion protein is less specific than the native SmD protein. Vice versa, only four of eight SmD sera reacted with the fusion protein. It should be noted however, that SmD was also expressed as a full-size β -galactosidase fusion protein in *E.coli* (Wagatsuma et al. 1993), but that this recombinant SmD antigen was not recognized by patient sera, although all sera recognized the natural Sm 16 kDa antigen on Western blot.

In conclusion, none of the described synthetic peptides nor the entire recombinant protein or parts of the molecule result in an immunoreactivity identical with the reactivity obtained with natural SmD.

It is an aim of the present invention to provide peptides which have a high reactivity for antibodies present in sera from patients with systemic lupus

erythematosus.

Another aim of the present invention is to provide methods for obtaining said peptides.

Another aim of the present invention is to provide methods of raising antibodies specifically reactive with peptides of said peptides, thereby mimicking said peptides.

Another aim of the present invention is to provide methods of raising anti-idiotypic antibodies specifically reactive with the afore mentioned antibodies.

Another aim of the present invention is to provide a pharmaceutical composition consisting of these peptides, for therapy or diagnosis.

Another aim of the present invention is to provide a diagnostic kit for systemic lupus erythematosus.

All these aims of the present invention are met by the following embodiments of the present invention.

According to its main embodiment the present invention relates to peptides containing less than 50 amino acids, comprising at least one dimer of the type XG, wherein X stands for a methylated arginine residue, and that are able to react with antibodies, with said methylation being crucial for the reaction between said peptide and said antibodies, and wherein said antibodies are present in sera from patients with systemic lupus erythematosus, or infectious, recurrent or chronic mononucleosis, or certain cancers which are related to infection with Epstein-Barr virus, such as Burkitt's lymphoma or nasopharyngeal carcinoma.

According to a further embodiment the present invention also relates to a peptide and/or chemical structure comprising any of the above mentioned peptides, fused to a linker molecule. The present invention also relates to peptides comprising and/or consisting of tandem repeats of at least two of any of the above mentioned peptides, or branched peptides that comprises at least one of the above mentioned peptides.

According to a more specific embodiment the present invention also relates to a method for producing any of the above mentioned peptides, by classical chemical synthesis, wherein methylated arginines are substituted for unmethylated

arginine residues at certain steps during the chemical synthesis. The present invention also relates to a method for producing any of the above mentioned peptides, wherein the primary amino acid sequence is produced by classical chemical synthesis, and wherein the arginine residues that precede glycine residues are subsequently methylated by contacting said peptides with a protein arginine methyltransferase. The present invention also relates to a method for producing any of the above mentioned peptides comprising the following steps:

5 (i) transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements such that said peptide or a protein comprising said peptide is expressed and/or secreted, (ii) culturing said transformed cellular host under conditions allowing expression of said protein or peptide and allowing a partial or optimal methylation of the arginines present in said peptide, and (iii) harvesting said peptide. The present invention also relates

10 to a method for producing any of the above mentioned peptides comprising the following steps: (i) transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements, such that said peptide or a protein comprising said peptide is expressed and/or secreted, (ii)

15 culturing said transformed cellular host under conditions allowing expression of said protein or said peptide, (iii) harvesting said protein or said peptide, and (iv) methylating arginine residues of said protein or said peptide by contacting with a protein arginine methyltransferase. According to a more specific embodiment the present invention also relates to any of the above mentioned methods, wherein

20 said host cell is a bacterial host or yeast or any other eukaryotic host cell which is preferably transformed with a recombinant baculovirus.

25

According to a preferred embodiment the present invention also relates to an antibody raised upon immunization with any of the above mentioned peptides, with said antibody being specifically reactive with the methylated forms of said peptide, and with said antibody being preferably a monoclonal antibody. The

30 present invention also relates to an anti-idiotypic antibody raised upon

immunization with any antibody as defined above, with said anti-idiotypic antibody being specifically reactive with said antibody, thereby mimicking the methylated form of any above mentioned peptide, and with said antibody being preferably a monoclonal antibody.

5 According to a more specific embodiment the present invention also relates to an immunotoxin molecule comprising and/or consisting of a cell recognition molecule being a peptide as defined above, or an antibody as defined above, covalently bound to a toxin molecule or active fragment thereof.

10 According to a further embodiment the present invention relates to any of the above mentioned peptides or antibodies or immunotoxine molecules or a composition thereof for use as a medicament. Said use can have the purpose of a medicament for treatment or of a diagnosticum for any of the following auto-immune diseases: systemic lupus erythematosus, discoid lupus erythematosus, scleroderma, dermatomyositis, rheumatoid arthritis, Sjögren's syndrome, or for
15 diseases in which Epstein-Barr virus can be implicated such as Burkitt's lymphoma or nasopharyngeal carcinoma, or infectious, recurrent or chronic mononucleosis. More specifically, the present invention relates to a treatment for auto-immune diseases by increasing the size of antigen-immune complexes, thereby improving the clearance of the formed immune complexes. The present invention also relates
20 to a treatment for auto-immune diseases by inducing a state of systemic hyporesponsiveness to the auto-antigen after oral administration of any of the above mentioned peptides or antibodies or immunotoxine molecules or a composition thereof, thereby preventing the pathogenic production of anti-self antibodies like anti-Sm antibodies or anti-DNA antibodies. The present invention
25 also relates to a diagnostic kit for use in detecting any of the afore mentioned diseases, wherein said kit comprises at least one of the above mentioned peptides or antibodies, and with said peptide or antibody being possibly bound to a solid support. More preferably said kit is comprising a range of said peptides or said antibodies, possibly in combination with native methylated SmD1 or SmD3 or
30 Sm69 and recombinant unmethylated SmD1 or SmD3 or Sm69, wherein said peptides are attached to specific locations on a solid substrate. More preferably

said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines. It has to be understood that certain peptides, or antibodies as defined above, alternatively, are not attached to a solid support but are provided in the binding solution to be used as competitors and/or to block other antibodies that are present in sera from patients with autoimmune diseases other than SLE, thereby decreasing or eliminating possible cross-reaction and/or aspecific binding.

We have demonstrated for the first time that well defined secondary modifications (mostly N^G,N^G -dimethylarginine) are present on the Arg residues of the C-terminal peptide, that are followed by a glycine residue. Moreover, we have raised evidence that the C-terminal peptide can only show an immunoreactivity almost identical to the immunoreactivity of natural SmD, if these arginine residues are methylated. These dimethylarginines present on the nine Arg positions of the C-terminus, have been demonstrated for the first time in the natural SmD1 molecule. In SmD2 no dimethylarginine was retrieved while in the C-terminus of SmD3 the four RG motifs in the C-terminus again were found to be dimethylated.

The amino acid N^G,N^G -dimethylarginine is the result of a post-translational modification which seems to occur predominantly in RNA binding proteins (Najbauer, 1993). These nuclear proteins are enzymatically modified by a nuclear protein methylase I (S-adenosyl-methionine: protein-arginine N-methyltransferase, E.C.2.1.1.23; Rajpurohit, et al., 1994). The structural specificity of this enzyme seems to be an arginine containing peptide with glycine in the C-flanking position as was shown by substrate evaluation with synthetic peptides (Rawal, 1995). Nevertheless, in the same study it was demonstrated that the entire molecule also plays an important though thus far unknown role in the methylation process. Interestingly, this cellular methylation process can be mimicked *in vitro* with purified methylase I as was illustrated with recombinant heterogeneous nuclear RNP protein A1 (Rajpurohit, et al. 1994)

From our results, we thus can conclude that in SmD immunoreactivity, at least 2 epitopes are involved. One of the epitopes is apparently present in the recombinant SmD1 molecule and can not be assigned to a linear epitope (epitope

mapping of *E. coli* recombinant SmD1, data not shown). This is in agreement with the discontinuous epitope described by Rokeach et al. (1992b). The epitope localized at the C-terminus both by epitope mapping with *E. coli* fusion fragments (Rokeach et al., 1992b; Rivkin et al., 1994) and synthetic peptides (Sabbatini et al., 1993a; James et al., 1994) could well be explained by the work of Rivkin et al. (1994). The latter group has demonstrated that a dipeptide repeat (Gly-Arg)₉ is recognized by 35% of sera from SLE patients but also by 15% of sera from other autoimmune diseases. This result is in contrast with the high SLE specificity of the anti-Sm antibodies. Indeed, the specificity of the unmodified C-terminal SmD1 peptide has not been thoroughly investigated by Rokeach (1992b) nor by James et al. (1994). Only Sabbatini (1993a) described a certain disease specificity for the C-terminal synthetic peptide. On one hand, Rivkin has shown that out of 32 sera positive for the (Gly-Arg)₉ peptide, only 6 sera are positive with native SmD. On the other hand, of the positive SmD sera identified on Western blot, only half of them are reactive with the unmodified C-terminal peptide (Rivkin: 4/8; Rokeach 9/19; Sabbatini: 5/9). Based on these results it can be concluded that immunoreactivity of the unmodified C-terminal peptide does not well correlate with natural SmD and is less SLE specific than natural SmD. In contrast, our results show that 15 out of 17 SmD positive sera are immunoreactive with the dimethylated C-terminal peptide while only one serum reacts with the unmodified C-terminal peptide. The 2 sera that do not recognize the dimethylated C-terminal peptide are immunoreactive with the total recombinant SmD and are apparently monospecific for the discontinuous epitope.

In conclusion, natural SmD1 contains nine dimethylated arginines at the C-terminus and this modification plays a crucial role in the SLE specific immunoreactivity of the SmD antigen.

According to its main embodiment the present invention relates to peptides that contain arginine residues that are immediately followed by a glycine residue, and wherein at least one arginine residue is methylated or dimethylated at one terminal aminogroup of the guanidino-group of the arginine residue, and wherein this methylation is a prerequisite for the peptide to be recognized by antibodies

that characterize certain diseases. Antibodies that are specifically reacting with this type of peptides can be found in sera from patients with systemic lupus erythematosus or related autoimmune diseases such as discoid lupus erythematosus, or patients with infectious mononucleosis, or recurrent or chronic mononucleosis, or that suffer from diseases in which Epstein-Barr virus has been implicated such as nasopharyngeal carcinoma and Burkitt's lymphoma.

Peptides are described which immunologically mimic the immunogenic determinants of self proteins recognized by the immune system in patients suffering from lupus erythematosus. A crucial aspect of such peptides is the fact that arginines followed by a glycine are methylated. One peptide (SmD1) has been demonstrated to contain a stretch of 9 consecutive arginine-glycine residues, wherein each arginine is methylated and wherein this methylation is necessary for specific recognition by antibodies present in sera of patients with lupus erythematosus. A second peptide (SmD3) has been demonstrated to contain isolated arginine-glycine residues, wherein the arginine is methylated. Also, a third peptide (Sm69) has been demonstrated to contain several domains characterised by several arginine-glycine residues, wherein the arginine is dimethylated. It is therefore anticipated that the presence of one dimethylated arginine, followed by a glycine can be sufficient for specific recognition by some antibodies present in sera of patients with lupus erythematosus. The invention therefore relates to those peptides wherein at least one arginine residue is followed by a glycine, wherein the arginine residue is methylated, and wherein this methylation is necessary for specific recognition by antibodies.

The term 'peptide' as used throughout the specification and claims refers to a polymer of amino acids and does not refer to a specific length of the product; thus, oligopeptides, polypeptides and proteins are included within the definition of 'peptide'. This term also does not refer to or exclude post-expression modifications of the peptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, peptides containing one or more analogues of an amino acid (including, for example, unnatural amino acids, PNA, etc.), polypeptides with substituted linkages, as well as other modifications

known in the art, both naturally occurring and non-naturally occurring.

Whenever the expression "peptide containing less than 50 amino acids" is used, this should be interpreted in a broad sense, as a means of circumscribing an essentially truncated version of entire immunoreactive proteins that still comprises the highly reactive domain characterized by the presence of methylated arginine residues. These peptides have a length of preferably 40, 30, 25, 20 or less amino acids. The present invention also relates to peptides having a length of 50, 60 or more amino acids without comprising the full length of the native protein. It is for practical purpose of peptide synthesis that peptides containing less than 50 amino acids are defined.

With 'immunogenic determinant' is meant, those chemical groupings comprising a primary amino acid sequence, and secondary modifications of the amino acid residues in a certain three-dimensional arrangement, that together determine the specific reactivity of the entire antigen for a raised antibody. Such antibody can also recognize different chemical groupings, which are then termed to 'immunologically mimic' the immunogenic determinant.

When secondary modifications of a peptide are said to be 'necessary' or 'crucial', or to 'be a prerequisite' for reacting with an antibody, the absence of said secondary modifications will result in a peptide of which the dissociation constant for interaction with said antibody will be at least two orders of magnitude higher than the dissociation constant for the interaction between said antibody and the peptide wherein the secondary modifications are present, preferably three orders of magnitude higher, and more preferably four orders of magnitude higher.

The term 'crossreaction' refers to the reaction of one antigen with antibodies developed against another antigen or antibodies that are found in sera from patients with different diseases.

The term 'methylated arginine' as used throughout the ensuing specification and claims is used in a broad sense and refers any methylated form. More preferably the term methylated refers to a dimethylated form of arginine in which one amino-group of the guanidino group of the arginine residue is substituted with one or two methyl groups. The term 'methylation' refers to the process that

results in said forms of arginine.

It has to be understood that the peptides of the present invention are not limited to include only those peptides that are recognized by antibodies specific for systemic lupus erythematosus, but also relate to those peptides that react with antibodies associated with related autoimmune diseases such as discoid lupus erythematosus and scleroderma or peptides that are related to infection with Epstein-Barr virus such as infectious, recurrent, or chronic mononucleosis. Several autoimmune diseases are related to systemic lupus erythematosus such as discoid lupus erythematosus, scleroderma. The antibodies that are present in sera of patients with these autoimmune diseases tend to recognize self- antigens, that are often localized in the nucleus and present in all tissues. As a result, similar or analogous epitopes are often recognized, thereby leading to crossreactivity, and difficult diagnosis.

The few antibodies present in sera from patients with systemic lupus erythematosus that are also able to react with recombinant and thus unmethylated C-terminal parts of SmD1, were shown to cross-react with the EBNA-1 Epstein-Barr virus associated nuclear antigen, of which the primary sequence displays homology with the primary sequence of SmD1 (Sabbatini et al., 1993b). EBNA-1 is the only protein which is consistently expressed during latency of Epstein-Barr virus. The extreme persistence of Epstein-Barr virus during latency can be explained by the very low immunogenicity of this viral protein. EBNA-1 most importantly contains six arginine residues which are followed by a glycine. It can therefore be anticipated that the frequency of finding specific antibodies in sera from patients with mononucleosis (infected with Epstein-Barr virus) or from people with recurrent, or chronic mononucleosis or with a history of mononucleosis, will increase, when antigens (such as EBNA-1) are used wherein the arginine residues that precede a glycine residue are methylated, as we have shown to be the case with the antibodies directed against SmD-1 in patients with lupus erythematosus.

According to a more specific embodiment the present invention relates to those peptides that contain methylated arginine residues that are preceding glycine residues, wherein said methylation is crucial for high-affinity interaction with

antibodies that are found in sera from patients with autoimmune diseases such as systemic lupus erythematosus, or that are infected with Epstein Barr virus.

The present invention also relates to those peptides wherein the arginine-glycine doublets are repeated, at least once, and more preferably 3 or 4 or 5 or 6 or 7 or 8 times, and even more preferably 9 times, as is the case with the natural antinuclear antigen SmD-1, and wherein at least one arginine residue that precedes a glycine residue is methylated, preferably dimethylated, with said methylation being necessary for specific reaction with antibodies, for instance with antibodies found in sera from patients with SLE.

In a more specific embodiment, the present invention relates to a peptide that is characterized by the amino acid sequence GRGRGRGRGRGRGRGRGRG (SEQ ID NO 16) wherein at least one and preferably each arginine is methylated, preferably dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the main immunogenic determinant of the C-terminal part of antinuclear antigen SmD1.

In a more specific embodiment, the present invention relates to a peptide that is characterized by the amino acid sequence DVEPKVKSKKREAVAGRG RGRGRGRGRGRGRGRGGPRR (SEQ ID NO 17) wherein at least one and preferably each arginine that precedes a glycine, is methylated, preferably dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the main immunogenic determinant of the C-terminal part of antinuclear antigen SmD1.

In a more specific embodiment, the present invention relates to a peptide that is characterized by the amino acid sequence ARGRGRGMGRG (SEQ ID NO 18) wherein at least one and preferably each arginine is methylated, preferably dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the main immunogenic determinant of the C-terminal part of antinuclear antigen SmD3.

In a more specific embodiment, the present invention also relates to a peptide that is characterized by the amino acid sequence KAQVAARGRGRGMGRGNIFQKRR (SEQ ID NO 19) wherein at least one and preferably each arginine that precedes a glycine is methylated, preferably

dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the main immunogenic determinant and its borders of the C-terminal part of antinuclear antigen SmD3.

According to a more specific embodiment, the present invention also relates to a peptide that comprises or consists of by the amino acid sequence GGQQDR GGRGRGGGGGYNRSSGGYEPRGRGGGRGGRMGGSDRGG (SEQ ID NO 20) or GGQQDRGGRGRGGGGGYN (SEQ ID NO 21) or SGGYEPRGRGGGRGGRMGGSDRGG (SEQ ID NO 22) or DFNRGGGNGRGGGRGRGG (SEQ ID NO 23) or DFNRGGGNGRGGGRGRGGPMGRGGYGGGGS (SEQ ID NO 24) or GDDRRGR GGYDRGGYRGRGGDRGGFRGGGRGGDRGGFG (SEQ ID NO 25) or GDDRRGRGGYDRGG (SEQ ID NO 26) or GGYRGRGGDRGGFRGGGRGGDRGGFG (SEQ ID NO 27) wherein at least one and preferably each arginine that precedes a glycine is methylated, preferably dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the main immunogenic determinant and its borders of the C-terminal part of antinuclear antigen Sm69.

According to a more specific embodiment, the present invention relates to a peptide that comprises or consists of the amino acid sequence DNHGRGRGRGRGGG (SEQ ID NO 28) or GGRGRGGSGGRGRGG (SEQ ID NO 29) or ERARGRGRGRGE (SEQ ID NO 30) wherein at least one and preferably each arginine that precedes a glycine is methylated, preferably dimethylated and even more preferably dimethylated in an asymmetric way, thereby mimicking the Epstein-Barr virus nuclear antigen1.

The present invention also relates to molecular structures in which at least part represents a peptide or antibody as defined above. Such molecular structures can result from fusion of peptides of the present invention with peptides and/or proteins and/or other molecules that are further characterized in that they -specifically interact with other peptides and/or proteins and/or molecular structures, enabling tagging and/or binding of the fused polypeptide and/or protein to specific tissue- or cell types or that allow for purification of said molecular structures due to the presence of for instance 4, or 5 or 6 consecutive histidine residues, or

-are cytotoxic to T-cells and/or B-cells such as cholera toxin, or
-allow for labelling by means of a radioactive or fluorescent or immunogold or enzymatic marker.

It may also be desirable in certain instances to join two or more peptides
5 together in one peptide structure, or to create branched peptides. One advantage of this arrangement is well known in the art and relates to diagnosis. When antibodies are used in an assay in order to detect the present antigens, tandem repeats or branched peptides of the antigens can increase the amount of immobilized antigens presented to the antibodies and thereby increase the
10 sensitivity of the assay. The sensitivity can be increased exponentially when the immobilized antigens are used together with a specific concentration of such antigens in a soluble form, thereby inducing the formation of crosslinked antigen-immunoprecipitates. A second advantage relates to therapy. The deposition of self-antigen autoimmune complexes in various tissues is an important step towards the
15 acquisition of a pathological condition. It is generally accepted that the main cause of said deposition is the insufficient blood clearance by the liver of the antigen-immune complexes due to the small size of said complexes. Administration of tandem repeats or branched forms of said peptides could increase the size of the formed antigen-immune complexes, and thereby increase the clearance and thus
20 decrease the deposition of said complexes.

The present invention also relates to circularized forms of said peptides, the advantage being well known in the art, and relating to an increased affinity of a conformationally constraint peptide as compared with the more randomly coiled forms of linear peptides.

25 In order to accommodate for eventual negative characteristics of the claimed peptides, such as rapid degradation, solubility, cytotoxic effects and so on, the skilled person will be able to design conservative as well as non-conservative amino acid substitutions, or substitutions with non-natural amino acids, PNA etc... These will generally account for less than 35 percent of a specific sequence. Such
30 peptides also include peptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally

occurring. It may be desirable in cases where the SmD peptides or other antigenic peptides of the present invention are highly polymorphic, to vary one or more of the amino acids so as to better mimic the different epitopes of several viral strains, or as recognized by antibodies in sera from patients with SLE or other autoimmune diseases.

The present invention also relates to any analogs of the peptides of the present invention.

The term "analog" as used throughout the specification or claims to describe the proteins or peptides of the present invention, includes any protein or peptide having an amino acid residue sequence substantially identical to a sequence specifically shown herein in which one or more residues have been conservatively substituted with a biologically equivalent residue. Examples of conservative substitutions include the substitution of hydrophobic residue such as isoleucine, valine, leucine or methionine for another, the substitution of one hydrophilic residue for another such as between arginine and lysine, between glutamine and asparagines, between glycine and serine, the substitution of one basic residue such as lysine, arginine or histidine for another, or the substitution of one acidic residue, such as aspartic acid or glutamic acid for another. Examples of allowable mutations according to the present invention can be found in Table 4.

Amino acids	Synonymous groups
Ser (S)	Ser, Thr, Gly, Asn
Arg (R)	Arg, His, Lys, Glu, Gln
Leu (L)	Leu; Ile, Met, Phe, Val, Tyr
Pro (P)	Pro, Ala, Thr, Gly
Thr (T)	Thr, Pro, Ser, Ala, Gly, His, Gln
Ala (A)	Ala, Pro, Gly, Thr
Val (V)	Val, Met, Ile, Tyr, Phe, Leu, Val
Gly (G)	Gly, Ala, Thr, Pro, Ser

Ile (I)	Ile, Met, Leu, Phe, Val, Ile, Tyr
Phe (F)	Phe, Met, Tyr, Ile, Leu, Trp, Val
Tyr (Y)	Tyr, Phe, Trp, Met, Ile, Val, Leu
Cys (C)	Cys, Ser, Thr, Met
His (H)	His, Gln, Arg, Lys, Glu, Thr
Gln (Q)	Gln, Glu, His, Lys, Asn, Thr, Arg
Asn (N)	Asn, Asp, Ser, Gln
Lys (K)	Lys, Arg, Glu, Gln, His
Asp (D)	Asp, Asn, Glu, Gln
Glu (E)	Glu, Gln, Asp, Lys, Asn, His, Arg
Met (M)	Met, Ile, Leu, Phe, Val

Table 4 Overview of the amino acid substitutions which could form the basis of analogs (muteins) as defined above.

The phrase "conservative substitution" also includes the use of a chemically derivatized residue in place of a non-derivatized residue provided that the resulting protein or peptide is biologically equivalent to the protein or peptide of the invention.

"Chemical derivative" refers to a protein or peptide having one or more residues chemically derivatized by reaction of a functional side group or peptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring. Examples of such derivatized molecules, include but are not limited to, those molecules in which free amino groups have been derivatized to form amine hydrochlorides, p-toluene sulfonyl groups, carbobenzoxy groups, t-butyloxycarbonyl groups, chloroacetyl groups or formyl groups. Free carboxyl groups may be derivatized to form salts, methyl and ethyl esters or other types of esters or hydrazides. Free hydroxyl groups may be derivatized to form O-acyl or O-alkyl derivatives. The imidazole nitrogen of histidine may be derivatized to form N-imbenzylhistidine. Also included as chemical derivatives are those proteins or peptides which contain one or more naturally-

occurring amino acid derivatives of the twenty standard amino acids. For examples : 4-hydroxyproline may be substituted for proline; 5-hydroxylysine may be substituted for lysine; 3-methylhistidine may be substituted for histidine; homoserine may be substituted for serine; and ornithine may be substituted for lysine. The peptides of the present invention also include any protein or peptide having one or more additions and/or deletions or residues relative to the sequence of a peptide whose sequence is shown herein, as long as the peptide is biologically equivalent to the proteins or peptides of the invention.

Furthermore, additional amino acids or chemical groups may be added to the amino- or carboxyl terminus for the purpose of creating a "linker arm" by which the peptide can conveniently be attached to a carrier. The linker arm will be at least one amino acid and may be as many as 60 amino acids but will most frequently be 1 to 10 amino acids. The nature of the attachment to a solid phase or carrier can be non-covalent as well as covalent. Possible arrangements of this nature are well described in the art. Natural amino acids such as histidine, cysteine, lysine, tyrosine, glutamic acid, or aspartic acid may be added to either the amino- or carboxyl terminus to provide functional groups for coupling to a solid phase or a carrier. However, other chemical groups such as, for example, biotin and thioglycolic acid, may be added to the termini which will endow the peptides with desired chemical or physical properties. The termini of the peptides may also be modified, for example, by N-terminal acetylation or terminal carboxy-amidation. In each instance, the peptide will preferably be as small as possible while still maintaining substantially all of the sensitivity of the larger peptide.

The peptides of the invention, and particularly the fragments, can be prepared by classical chemical synthesis. The synthesis can be carried out in homogeneous solution or in solid phase. For instance, the synthesis technique in homogeneous solution which can be used is the one described by Houbenweyl in the book entitled "Methode der organischen chemie" (Method of organic chemistry) edited by E. Wunsh, vol. 15-I et II. THIEME, Stuttgart 1974. The polypeptides of the invention can also be prepared in solid phase according to the methods described by Atherton and Shepard in their book entitled "Solid phase

peptide synthesis" (IRL Press, Oxford, 1989). The methylated forms of the claimed peptides can be obtained by substituting the methylated arginine derivatives for the normal arginine derivatives during the classical chemical synthesis, or by contacting the unmethylated peptides after synthesis with a protein arginine methyl transferase enzyme of any eukaryotic origin.

The polypeptides according to this invention can also be prepared by means of recombinant DNA techniques as described by Maniatis et al., Molecular Cloning: A Laboratory Manual, New York, Cold Spring Harbor Laboratory, 1982) by insertion of a polynucleic acid sequence encoding the claimed peptides or part of the claimed peptides in an appropriate vector and transforming a suitable host with said vector. This recombinant expression vector comprises a polynucleic acid or a part thereof as defined above, operably linked to prokaryotic, eukaryotic or viral transcription and translation control elements. In addition this sequence can be operably linked with sequences that allow for secretion of the claimed peptides. The term 'vector' may comprise a plasmid, a cosmid, a phage or a virus or a transgenic organism. Particularly useful may be BCG or adenoviral vectors, as well as avipox recombinant viruses.

The recombinant peptides can be methylated *in vitro*, by contacting the expressed and/or secreted peptides with a protein arginine methyl transferase of any eukaryotic origin, or *in vivo* by choosing the appropriate host, like yeast, or any eukaryotic cell, and more preferably by using the baculovirus transformation system.

The present invention does not exclude the option of using additional proteins like the BTG1 and TIS21 proteins which have been demonstrated to be essential for methylation *in vivo* (as a co-expressed protein) or to be required for optimal methylation *in vitro* (Lin et al., 1996), or any other proteins, peptides or chemical substances that can optimize the level of methylation.

Also any of the known purification methods for recombinant peptides can be used for the production of the recombinant peptides of the present invention.

The present invention also relates to a recombinant expression vector comprising a polynucleic acid or a part thereof as defined above, operably linked

to prokaryotic, eukaryotic or viral transcription and translation control elements.

In general, said recombinant vector will comprise a vector sequence, an appropriate prokaryotic, eukaryotic or viral promoter sequence followed by a nucleotide sequence encoding a peptide as defined above, with said recombinant vector allowing the expression and/or secretion of any one of the polypeptides as defined above in a prokaryotic, or eukaryotic host or in living mammals when injected as naked DNA.

Also any of the known purification methods for recombinant proteins may be used for the production of the recombinant polypeptides of the present invention.

The term "vector" may comprise a plasmid, a cosmid, a phage, or a virus or a transgenic animal. Particularly useful for vaccine development may be BCG or adenoviral vectors, as well as avipox recombinant viruses.

The present invention also relates to a method for the production of a recombinant polypeptide as defined above, comprising:

- transformation of an appropriate cellular host with a recombinant vector, in which a polynucleic acid or a part thereof according to as defined above has been inserted under the control of appropriate regulatory elements,
- culturing said transformed cellular host under conditions enabling the expression and/or secretion of said insert, and,
- harvesting said polypeptide.

The term "recombinantly expressed" used within the context of the present invention refers to the fact that the proteins of the present invention are produced by recombinant expression methods be it in prokaryotes, or lower or higher eukaryotes as discussed in detail below.

The term "lower eukaryote" refers to host cells such as yeast, fungi and the like. Lower eukaryotes are generally (but not necessarily) unicellular. Preferred lower eukaryotes are yeasts, particularly species within Saccharomyces, Schizosaccharomyces, Kluveromyces, Pichia (e.g. Pichia pastoris), Hansenula (e.g. Hansenula polymorpha), Yarrowia, Schwaniomyces, Schizosaccharomyces, Zygosaccharomyces and the like. Saccharomyces cerevisiae, S. carlsbergensis and

K. lactis are the most commonly used yeast hosts, and are convenient fungal hosts.

The term "prokaryotes" refers to hosts such as E.coli, Lactobacillus, Lactococcus, Salmonella, Streptococcus, Bacillus subtilis or Streptomyces. Also these hosts are contemplated within the present invention.

The term "higher eukaryote" refers to host cells derived from higher animals, such as mammals, reptiles, insects, and the like. Presently preferred higher eukaryote host cells are derived from Chinese hamster (e.g. CHO), monkey (e.g. COS and Vero cells), baby hamster kidney (BHK), pig kidney (PK15), rabbit kidney 13 cells (RK13), the human osteosarcoma cell line 143 B, the human cell line HeLa and human hepatoma cell lines like Hep G2, and insect cell lines (e.g. Spodoptera frugiperda). The host cells may be provided in suspension or flask cultures, tissue cultures, organ cultures and the like. Alternatively the host cells may also be transgenic animals.

The term "recombinant polynucleotide" or "nucleic acid" intends a polynucleotide or nucleic acid of genomic, cDNA, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation : (1) is not associated with all or a portion of a polynucleotide with which it is associated in nature, (2) is linked to a polynucleotide other than that to which it is linked in nature, or (3) does not occur in nature.

The term "recombinant host cells", "host cells", "cells", "cell lines", "cell cultures", and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which can be or have been, used as recipients for a recombinant vector or other transfer polynucleotide, and include the progeny of the original cell which has been transfected. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

The term "replicon" is any genetic element, e.g., a plasmid, a chromosome, a virus, a cosmid, etc., that behaves as an autonomous unit of polynucleotide replication within a cell; i.e., capable of replication under its own control.

The term "vector" is a replicon further comprising sequences providing replication and/or expression of a desired open reading frame.

The term "control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, splicing sites and terminators; in eukaryotes, generally, such control sequences include promoters, splicing sites, terminators and, in some instances, enhancers. The term "control sequences" is intended to include, at a minimum, all components whose presence is necessary for expression, and may also include additional components whose presence is advantageous, for example, leader sequences which govern secretion.

The term "promoter" is a nucleotide sequence which is comprised of consensus sequences which allow the binding of RNA polymerase to the DNA template in a manner such that mRNA production initiates at the normal transcription initiation site for the adjacent structural gene.

The expression "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences.

The polynucleic acids encoding the peptides of the present invention and inserted into the vector sequence may be attached to a signal sequence. Said signal sequence may be that from any source, e.g. the IgG or tissue plasminogen activator (tpa) leader sequence for expression in mammalian cells, or the α -mating factor sequence for expression into yeast cells.

A variety of vectors may be used to obtain the peptides of the present invention. Lower eukaryotes such as yeasts and glycosylation mutant strains are typically transformed with plasmids, or are transformed with a recombinant virus. The vectors may replicate within the host independently, or may integrate into the host cell genome.

Higher eukaryotes may be transformed with vectors, or may be infected with a recombinant virus, for example a recombinant vaccinia virus. Techniques and vectors for the insertion of foreign DNA into vaccinia virus are well known in the art, and utilize, for example homologous recombination. A wide variety of viral promoter sequences, possibly terminator sequences and poly(A)-addition sequences, possibly enhancer sequences and possibly amplification sequences, all required for the mammalian expression, are available in the art. Vaccinia is particularly preferred since vaccinia halts the expression of host cell proteins. Vaccinia is also very much preferred since it allows the expression of f.i. peptides of the present invention in cells or individuals which are immunized with the live recombinant vaccinia virus. For vaccination of humans the avipox and Ankara Modified Virus (AMV) are particularly useful vectors.

Also known are insect expression transfer vectors derived from baculovirus Autographa californica nuclear polyhedrosis virus (AcNPV), which is a helper-independent viral expression vector. Expression vectors derived from this system usually use the strong viral polyhedrin gene promoter to drive the expression of heterologous genes. Different vectors as well as methods for the introduction of heterologous DNA into the desired site of baculovirus are available to the man skilled in the art for baculovirus expression. Also different signals for posttranslational modification recognized by insect cells are known in the art.

The present invention also relates to a host cell transformed with a recombinant vector as defined above.

The present invention also relates to antibodies that are specifically raised against the peptides of the present invention, preferably against those peptides wherein the arginines that precede a glycine residue are methylated. These antibodies may be polyclonal or monoclonal. To prepare antibodies a host animal is immunized using the peptides of the present invention in a pharmaceutically acceptable carrier, wherein at least one of the arginines that precede a glycine residue of said peptides is methylated. Pharmaceutically acceptable carriers include any carrier that does not itself induce the production of antibodies harmful to the

individual receiving the composition. Suitable carriers are typically large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers; and inactive virus particles. Such carriers are well known to those of ordinary skill in the art.

5 Preferred adjuvants to enhance effectiveness of the composition include, but are not limited to : aluminum hydroxide (alum), N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP) as found in U.S. Patent No. 4,606,918, N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE) and RIBI, which contains three components extracted from
10 bacteria, monophosphoryl lipid A, trehalose dimycolate, and cell wall skeleton (MPL+TDM+CWS) in a 2% squalene/Tween 80 emulsion. Any of the 3 components MPL, TDM or CWS may also be used alone or combined 2 by 2. Additionally, adjuvants such as Stimulon (Cambridge Bioscience, Worcester, MA) or SAF-1 (Syntex) may be used. Further, Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA) may be used for non-human applications and research purposes.

The immunogenic compositions typically will contain pharmaceutically acceptable vehicles, such as water, saline, glycerol, ethanol, etc. Additionally,
15 auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, preservatives, and the like, may be included in such vehicles.

Typically, the immunogenic compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared. The
20 preparation also may be emulsified or encapsulated in liposomes for enhanced adjuvant effect. The proteins may also be incorporated into Immune Stimulating Complexes together with saponins, for example Quil A (ISCOMS).

Immunogenic compositions used to raise antibodies comprise a 'sufficient amount' or 'an immunologically effective amount' of the peptides of the present
25 invention, as well as any other of the above mentioned components, as needed. 'Immunologically effective amount', means that the administration of that amount

to an individual, either in a single dose or as part of a series, is effective to provoke an immune response and to raise antibodies, as defined above. This amount varies depending upon the health and physical condition of the individual, the taxonomic group of the individual to be treated (e.g. nonhuman primate, primate, rabbit, etc.), the capacity of the individual's immune system to synthesize antibodies, the immunogenicity of the antigenic peptide, and its mode of administration, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials. Usually, the amount will vary from 0.01 to 1000 $\mu\text{g}/\text{dose}$, more particularly from 0.1 to 100 $\mu\text{g}/\text{dose}$.

The immunogenic compositions are conventionally administered parenterally, typically by injection, for example, subcutaneously or intramuscularly. Additional formulations suitable for other methods of administration include oral formulations and suppositories. Dosage treatment may be a single dose schedule or a multiple dose schedule. The vaccine may be administered in conjunction with other immunoregulatory agents.

The host serum or plasma is collected following an appropriate time interval to provide a composition comprising antibodies reactive with the peptides of the present invention. The gamma globulin fraction or the IgG antibodies can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other techniques known to those skilled in the art. The antibodies are substantially free of many of the adverse side effects which may be associated with other anti-viral agents such as drugs, for the treatment of infectious, chronic, or recurrent mononucleosis. Such antibodies may also be used to diagnose certain diseases, such as Burkitt's lymphoma, wherein Epstein-Barr virus has been implicated.

The term 'immunogenic' refers to the ability of a substance to cause a humoral and/or cellular response, whether alone or when linked to a carrier, in the presence or absence of an adjuvant.

The antibodies of the claimed invention may also be monoclonals that are prepared with said antibody being specifically reactive with any of said peptides,

and with said antibody being preferably a monoclonal antibody.

The monoclonal antibodies of the invention can be produced by any hybridoma liable to be formed according to classical methods from splenic cells of an animal, particularly from a mouse or rat, immunized against the claimed peptides of the present invention on the one hand, and of cells of a myeloma cell line on the other hand, and to be selected by the ability of the hybridoma to produce the monoclonal antibodies recognizing the methylated forms of the peptides which has been initially used for the immunization of the animals.

The antibodies involved in the invention can be labelled by an appropriate label of the enzymatic, fluorescent, or radioactive type.

The monoclonal antibodies according to this preferred embodiment of the invention may be humanized versions of mouse monoclonal antibodies made by means of recombinant DNA technology, departing from parts of mouse and/or human genomic DNA sequences coding for H and L chains or from cDNA clones coding for H and L chains.

Alternatively the monoclonal antibodies according to this preferred embodiment of the invention may be human monoclonal antibodies. These antibodies according to the present embodiment of the invention can also be derived from human peripheral blood lymphocytes of patients with SLE or any other autoimmune disease or with infectious, or recurrent or chronic mononucleosis. Such human monoclonal antibodies are prepared, for instance, by means of human peripheral blood lymphocytes (PBL) repopulation of severe combined immune deficiency (SCID) mice (for recent review, see Duchosal et al. 1992) or by screening Epstein Barr-virus-transformed lymphocytes of infected or vaccinated individuals for the presence of reactive B-cells by means of the antigens of the present invention.

The present invention also relates to the anti-idiotypic antibodies that are raised upon immunization with an antibody as defined above and that specifically react with said antibodies, thereby mimicking the peptides of the present invention. The methods for production of monoclonal anti-idiotypic antibodies, which are well known in the art, have been described, for instance, by Gheuens

et MacFarlin (1982).

The present invention also relates to truncated versions or single chain versions of the antibodies and anti-idiotypic antibodies as defined above, that have retained their original specificity for reacting with the antigens.

5 The present invention also relates to proteins or peptides that mimic the antibodies as defined above such as microproteins as can be obtained by phage display or the highly variable domain of a recombinant antibody as obtained by screening upon repertoire cloning.

10 The present invention also relates to a method for detecting antibodies that specifically react with the peptides or anti-idiotypic antibodies of the present invention, present in a biological sample, comprising:

- 15 (i) contacting the biological sample to be analysed for the presence of said antibodies with a peptide or anti-idiotypic antibody as defined above,
(ii) detecting the immunological complex formed between said antibodies and said peptide or anti-idiotypic antibody.

20 The present invention also relates to a reverse method for detecting the peptides and/or the anti-idiotypic antibodies of the present invention with antibodies present in a biological sample that specifically react with methylated forms of said peptides and/or anti-idiotypic antibodies that mimic such peptides, comprising:

- (i) contacting the biological sample to be analysed for the presence of said peptides or anti-idiotypic antibodies with the antibodies as defined above,
(ii) detecting the immunological complex formed between said antibodies and said peptide or anti-idiotypic antibody.

25 The methods as defined above, can be used in the diagnosis of autoimmune diseases such as systemic lupus erythematosus, discoid lupus erythematosus, scleroderma, dermatomyositis, rheumatoid arthritis, Sjögren's syndrome, or diseases in which Epstein-Barr virus can be implicated such as infectious, recurrent or chronic mononucleosis, or Burkitt's lymphoma, or nasopharyngeal carcinoma,
30 or Hodgkin's disease, or of certain cancers such as Ewing sarcoma, or malignant melanoma of soft tissue.

According to a specific embodiment, the present invention relates to the development of a diagnostic technique that allows differentiation between those autoimmune diseases in which the characteristic antibodies often crossreact with the same antigen, thus resulting in difficult and slow diagnosis. Such diagnostic technique can be obtained by the simultaneous use of several antigens, methylated and unmethylated, and at least two epitopes, a methylated and a non-methylated form of any of the claimed peptides and/or anti-idiotypic antibodies of the present invention.

The present invention also relates to a diagnostic kit for use in detecting the presence of said antibodies, said kit comprising at least one peptide or anti-idiotypic antibody or microprotein as defined above, with said peptide or anti-idiotypic antibody or microprotein being preferably bound to a solid support.

The present invention also relates to a diagnostic kit for determining the type of autoimmune disease or the type of infection or to characterize certain cancers, said kit comprising at least one peptide or anti-idiotypic antibody or microprotein as defined above, with said peptide or anti-idiotypic antibody or microprotein being preferably bound to a solid support.

The present invention also relates to a diagnostic kit as defined above, said kit comprising a range of said peptides and/or anti-idiotypic antibodies or microprotein which are attached to specific locations on a solid substrate.

The present invention also relates to a diagnostic kit as defined above, wherein said solid support is a membrane strip and said peptides and/or anti-idiotypic antibodies or microproteins are coupled to the membrane in the form of parallel lines.

The immunoassay methods according to the present invention may utilize for instance single or specific oligomeric antigens, dimeric antigens, as well as combinations of single or specific oligomeric antigens. The peptides of the present invention may be employed in virtually any assay format that employs a known antigen to detect antibodies that characterize a certain disease or infection. A common feature of all of these assays is that the antigenic peptide or anti-idiotypic antibody or microprotein is contacted with the body component suspected of

containing the antibodies under conditions that permit the antigen to bind to any such antibody present in the component. Such conditions will typically be physiologic temperature, pH and ionic strength using an excess of antigen. The incubation of the antigen with the specimen is followed by detection of immune complexes comprised of the antigen.

Design of the immunoassays is subject to a great deal of variation, and many formats are known in the art. Protocols may, for example, use solid supports, or immunoprecipitation. Most assays involve the use of labelled antibody or peptide; the labels may be, for example, enzymatic, fluorescent, chemiluminescent, radioactive, or dye molecules. Assays which amplify the signals from the immune complex are also known; examples of which are assays which utilize biotin and avidin or streptavidin, and enzyme-labelled and mediated immunoassays, such as ELISA assays.

The immunoassay may be, without limitation, in a heterogeneous or in a homogeneous format, and of a standard or competitive type. In a heterogeneous format, the peptide or anti-idiotypic antibody or microprotein is typically bound to a solid matrix or support to facilitate separation of the sample from the peptide or anti-idiotypic antibody or microprotein after incubation. Examples of solid supports that can be used are nitrocellulose (e.g., in membrane or microtiter well form), polyvinyl chloride (e.g., in sheets or microtiter wells), polystyrene latex (e.g., in beads or microtiter plates, polyvinylidene fluoride (known as Immunolon™), diazotized paper, nylon membranes, activated beads, and Protein A beads. For example, Dynatech Immunolon™ 1 or Immunolon™ 2 microtiter plates or 0.25 inch polystyrene beads (Precision Plastic Ball) can be used in the heterogeneous format. The solid support containing the antigenic peptides or anti-idiotypic antibodies or microprotein is typically washed after separating it from the test sample, and prior to detection of bound antibodies. Both standard and competitive formats are known in the art.

In a homogeneous format, the test sample is incubated with the combination of antigens in solution. For example, it may be under conditions that will precipitate any antigen-antibody or anti-idiotypic antibody-antibody or microprotein-

antibody complexes which are formed. Both standard and competitive formats for these assays are known in the art. For instance, to characterize SLE or systemic lupus erythematosus in a standard format, the amount of SLE-antibodies in the antibody-antigen complexes is directly monitored. This may be accomplished by determining whether a second type of labelled anti-xenogenetic (e.g. anti-human) antibodies which recognize an epitope on the first type of SLE-antibodies will bind due to complex formation. In a competitive format, the amount of SLE-antibodies in the sample is deduced by monitoring the competitive effect on the binding of a known amount of labelled antibody (or other competing ligand) in the complex. The detection of SLE-antibodies for diagnosis of SLE is used as an illustration. Wherever the term "SLE-antibodies" is used throughout the specification, this should not be considered as limitative. Like wise, the other autoimmune diseases are diagnosed by detection of other antibodies, and mononucleosis is diagnosed by detection of anti-Epstein-Barr virus antibodies.

Complexes formed comprising SLE-antibody (or in the case of competitive assays, the amount of competing antibody) are detected by any of a number of known techniques, depending on the format. For example, unlabelled SLE-antibodies in the complex may be detected using a conjugate of anti-xenogenetic Ig complexed with a label (e.g. an enzyme label).

In an immunoprecipitation or agglutination assay format the reaction between the SLE-antigens and the SLE-antibody forms a network that precipitates from the solution or suspension and forms a visible layer or film of precipitate. If no SLE-antibody is present in the test specimen, no visible precipitate is formed.

Currently, there exist three specific types of particle agglutination (PA) assays. These assays are used for the detection of antibodies to various antigens when coated to a support. One type of this assay is the hemagglutination assay using red blood cells (RBCs) that are sensitized by passively adsorbing antigen (or antibody) to the RBC. The addition of specific antigen antibodies present in the body component, if any, causes the RBCs coated with the purified antigen to agglutinate.

To eliminate potential non-specific reactions in the hemagglutination assay,

two artificial carriers may be used instead of RBC in the PA. The most common of these are latex particles. However, gelatin particles may also be used. The assays utilizing either of these carriers are based on passive agglutination of the particles coated with purified antigens.

5 The antigenic peptides of the present invention will typically be packaged in the form of a kit for use in these immunoassays. The kit will normally contain in separate containers the antigenic peptide or anti-idiotypic antibody, control antibody formulations (positive and/or negative), labelled antibody when the assay format requires the same and signal generating reagents (e.g. enzyme substrate)
10 if the label does not generate a signal directly. The antigenic peptide or anti-idiotypic antibody may be already bound to a solid matrix or separate with reagents for binding it to the matrix. Instructions (e.g. written, tape, CD-ROM, etc.) for carrying out the assay usually will be included in the kit.

The solid phase selected can include polymeric or glass beads, nitrocellulose, microparticles, microwells of a reaction tray, test tubes and magnetic beads. The signal generating compound can include an enzyme, a luminescent compound, a chromogen, a radioactive element and a chemiluminescent compound. Examples of enzymes include alkaline phosphatase, horseradish peroxidase and beta-galactosidase. Examples of enhancer compounds
20 include biotin, anti-biotin and avidin. Examples of enhancer compounds binding members include biotin, anti-biotin and avidin. In order to block the effects of rheumatoid factor-like substances, the test sample is subjected to conditions sufficient to block the effect of rheumatoid factor-like substances. These conditions comprise contacting the test sample with a quantity of for instance anti-
25 human IgG to form a mixture, and incubating the mixture for a time and under conditions sufficient to form a reaction mixture product substantially free of rheumatoid factor-like substance.

The present invention particularly relates to an immunoassay format in which several peptides of the invention are coupled to a membrane in the form of
30 parallel lines. This assay format is particularly advantageous for allowing a discrimination between the separate autoimmune diseases. The antigens that are

immobilized on the membrane will preferentially be the methylated and unmethylated form of poly(Arg-Gly), combined with native and thus methylated SmD1 and/or SmD3 and/or Sm69, and unmethylated, recombinant SmD1 and/or SmD3 and/or Sm69.

5

Figure Legends

Figure 1: HPLC profile of the Endo-Lys digest.

Figure 2: Immunodot of HPLC fractions with 5 patients sera and 1 control serum.

Figure 3: Immunodot of the C-terminal peptide (C-term mod) and without (C-term nt mod) dimethylarginine, and of the recombinant (baculo SmD, coli SmD) and natural protein (native). Strips were incubated with a anti-SmD positive serum (+) and a control serum (-). Total protein staining (Aurodyne) was performed on the third strip.

Figure 4: LIA with modified (dimethyl arginine) C terminal peptide (fraction 15 from EndoLys-C digest, line 1 on the strip), and non-modified C terminal peptide (fraction 8 from the EndoLys-C digest, line 2 on the strip), both applied in equal amounts (60 ng). Additionally, 7, 15 and 30 ng of recombinant SmD1 from baculovirus- or E. Coli-infected insect cells (resp. 4,5,6 and 7,8,9) as well as 15 and 30 ng of a mixture of gel-purified SmD (native) were applied to the strips. The total protein staining (Aurodyne) was performed on the first strip. The strips were incubated with (A) a panel of anti-SmD positive sera selected by INNO-LIA ANA from ANF-positive sera, (B) a panel of anti-SmD positive sera selected by INNO-LIA ANA from a cohort of SLE patients diagnosed according to the ACR criteria, (C) sera selected from MCTD patients (control panel) and (D) sera selected from ANF-negative sera (control panel). No reactivity was observed with sera from the control panels.

Examples

Example 1. Sera

Sm positive sera were obtained from the Department of Rheumatology of the University clinic in Ghent, Belgium (Dr. De Keyser and Dr. Veys). These sera were identified by microgel diffusion blotting (MDB) using rabbit thymus extract (Zeus, Bayer, Raritan, USA) as substrate (De Keyser *et al.*, 1990). The Sm-positivity was defined by a positive immunoreaction at the same molecular weight position (approx. 14 kDa) as the α -SmD reference serum.

Example 2. Isolation of native SmD

The snRNP particles are purified from HeLa nuclear extracts by immuno affinity chromatography (R. Lührmann, Marburg, Germany).

The snRNP particles are received in 20 mM Hepes/KOH, pH 7.9 - 250 à 420 mM NaCl - 5% glycerol - 1.5M $MgCl_2$ - 0.2 mM EDTA - 0.5 mM DTE - 0.5 mM PMSF. SmD is isolated from these particles as described by Lehmeier *et al.* (1990) with some modifications. Briefly, in a first step, snRNPs are concentrated in a centricon concentrator (30K centriep, Amicon) to a final volume of 5-10 mg/ml. Subsequently, the snRNPs are dissolved in Laemmli sample buffer, separated in a preparative 15% Laemmli gel (1 cm thick, 14 cm well; protein load 3mg) and stained with Coomassie Brilliant Blue.

The 14 kDa SmD (containing SmD1, SmD2 and SmD3) band is cut from the gel, rinsed in water and cut into 1 mm³ cubes. Proteins are eluted from the polyacrylamide gel in the BioRad apparatus according to the manufacture's instructions. Residual SDS and CoomassieBB are removed from the electroeluted

proteins by ion-pair extraction (precipitation of the dried protein with acetone/acidic acid/triethylamine/water:85/5/5/5). The pellet is dissolved in 6M ureum, 0.1 M acetic acid (glacial) and immediately neutralized with 1.5 M Tris-HCl.

Protein concentration is determined by MicroBCA method (Pierce,USA), an average yield of 80 μ g SmD/mg snRNPs is obtained.

Example 3. Expression of SmD1 as short mTNF-fusion in *E. coli* and purification of the fusion protein.

The SmD1 coding sequence (357 bp) was isolated from a cDNA clone bought from Organon Technika as a 367 bp PCR fragment by using *pfu* polymerase (T_m : 55°C). This PCR fragment was cut with BamHI and XbaI and inserted into the BamHI/XbaI cut expression vector pIGFH111. This expression vector was transformed to *E.coli* expression strain SG4044(pcl857). Induction of this vector/strain combination at 37°C showed a strong signal of ± 18 kDa on CBB stained gels and on Western blot. Upon localisation analysis the protein proved to be present in the soluble fraction. No significant proteolytic breakdown could be observed. Bacterial cells derived from three litre culture were suspended in lysis buffer (10 mM Tris- 100 mM KCl pH 6.8) until 3 times the amount of wet cells. Prior to lysis by French press, ϵ -aminocaproic acid, DTT, and PMSF were added to a final concentration of 25 mM, 1 mM and 2mM respectively. The cell suspension was forced twice through the French press and pressure was kept at 14000 psi. Before centrifugation, the lysate was diluted with lysis buffer (5 times the wet cell weight) and was centrifuged for 20 minutes at 27,000 g at 4°C. Guanidine HCl was added to the supernatant to an end concentration of 4.5 M. The recombinant fusion protein, containing a His-tag, was purified in a single step by metal affinity chromatography (Ni-IMAC sepharose). Chromatography was performed at room temperature. The column was loaded with 1 column volume of NiCl_2 (5mg/ml), washed with water and equilibrated with buffer A (6 M guanidine HCl, 0.1 M sodium phosphate, 0.05% TritonX100, pH 6.5) . The proteins were

loaded on Ni^{2+} chelating sepharose (Pharmacia, Sweden; approx. 18 mg protein/ml gel) and the column was washed with 4 bed volumes of buffer A. SmD1 was eluted with a linear gradient of buffer B (6 M guanidine HCl, 0.1 M sodium phosphate, 0.05% TritonX100, pH 3.5) and the protein eluted between 70% and 90% buffer B.

Example 4. Expression of SmD1 as short mTNF-fusion in the baculoviral system and purification of the fusion protein

The cDNA gene coding for the mTNF-His6-hSmD fusion protein was isolated from the bacterial expression plasmid pIGFH111hSmD (see example 3) as a 520 bp *Dra*I-*Xba*I fragment, and inserted in the *Bam*HI (filled in)-*Xba*I opened baculo transfer plasmid pVL1393, resulting in the recombinant transfer plasmid pVLTNFH6hSmD (see Fig2). The fusion gene is here under transcriptional control of the baculovirus strong polyhedrin promoter. The pVmTNFH6hSmD1 baculo transfer vector was used to generate recombinant mTNF-His6-hSmD1 baculovirus following the baculogold transfection approach (PharMingen, San Diego, USA). Infection of *Spodoptera frugiperda* cells (Sf9) with the recombinant virus resulted in the expression of a 18 kDa protein which was recognized on Western blot by a monoclonal antibody specific for SmD (Progen, Heidelberg, Germany, data not shown). Using the cell lysate for testing the specificity of different human sera was not feasible since a high aspecific background reaction of the human sera with baculoviral proteins masked possible specific SmD recognition. The SmD fusion protein was therefore purified by Ni-IMAC purification as described previously with one adaptation: following french press, the cell lysate was precipitated and redissolved in buffer A (see example 3).

Example 5. Sequence and mass analysis of natural, *E. coli*, and baculoviral SmD1

Natural SmD electroeluted from HeLa nuclear extracts immobilized on a PVDF

membrane in a ProSpin device (Perkin Elmer, California, USA) was subjected to endoLys-C digestion to obtain detailed sequencing data of internal peptides. The membrane was incubated with 100 mM Tris pH 8.2, 1% hydrogenated TritonX-100, 1 mM K₃-EDTA, 10% acetonitrile and 0.5 µg enzyme. The digestion was performed overnight at 37°C. The peptide mixture was separated on a C4 Vydac HPLC-column (using a gradient of 10-70% solvent B: 70% acetonitrile/0.1% TFA) and a flow rate of 0.2 ml/min. The eluted peptide peaks were manually recovered. In the C-terminal 25-mer peptide of SmD1 nine dimethylarginine residues were sequenced, only the last two arginines were unmodified. The position of N^G, N^G-dimethylarginine in the sequence chromatogram was confirmed by applying the pure modified amino acid (Sigma, St Louis, USA) as standard. This modification was absent in recombinant SmD1 from *E. coli* (revealed in the course of sequencing peptides generated by endo-GluC) implying that the modification, resulting from the action of methyltransferase, does not occur in *E. coli*.

This conclusion was confirmed by mass analysis of *E. coli* recombinant SmD1. This protein, eluting in a single peak upon reversed-phase chromatography, was analysed by electrospray on a Bio-Q quadrupole mass spectrometer equipped with an electrospray ion source (Fisons). Ten µl of the sample solution containing 20 pmol in 50% acetonitrile-1% acetic acid was analysed. Calibration of the scans was performed with 50 pmol horse heart myoglobin. The sample contained 3 masses: 17,435 Da, 17,305 Da, and 16,992 Da corresponding respectively to the full size protein, the protein without the N-terminal methionine, and the protein lacking the N-terminal Met and the C-terminal Arg-Arg. From these results, it can be concluded that the purified *E. coli* recombinant SmD1 is the intact, unmodified molecule and that the lack of specific immunoreactivity of the recombinant SmD1 is not due to loss of the C-terminus.

Mass analysis of baculoviral recombinant SmD1 showed a heterogeneous result: one of the major mass peaks (17,297 Da) could be assigned to the unmodified protein lacking the N-terminal methionine while within the minor peaks masses of 17,629 and 17,711 could be tentatively assigned to the presence of 7 and 10 dimethylarginines.

Example 6. Epitope mapping of baculo SmD1

Baculo SmD1 fusion protein was digested with EndoGlu-C as follows: 300 μ g TCA-precipitated protein was dissolved in 50 μ l 100 mM NH_4 -acetate buffer pH 4.3. The EndoGlu-C enzyme (Boehringer, Mannheim, Germany) was added at a ratio of 1/100 and the mixture was incubated overnight at 26°C. The digest was subsequently vacuum dried (SpeedVac), redissolved in 0.1 % TFA -20% acetic acid and the peptides were separated on a reversed-phase HPLC column (C_4 -Vydac). Peptide peaks were manually recovered. A similar approach was followed for EndoLys-C (Boehringer, Mannheim, Germany) digestion of baculo SmD1 with the following modifications: the protein is dissolved in 50 μ l 100 mM Tris-HCl, pH 8, 10% acetonitrile, 10 mM K_3EDTA , and enzyme is added at a ratio of 1/120.

The HPLC fractions were vacuum dried and dissolved in 10% acetonitrile, 50 mM carbonate buffer pH 9.6. From each fraction 2 μ l was spotted on ABC nylon membrane (Pall, NY). After spotting, the membranes were blocked for 1 hour in 0.5% caseine in PBS to which 0.1% 0.25 glycine is added. Subsequently, the membranes are incubated overnight with serum (1/100) in 0.5% caseine in PBS supplemented with Triton X705 and 2.03 g/L $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$. The membranes were washed 3 times for 3 min. in PBS, 0.05% Tween20 and incubated with anti-human IgG (1/8000) conjugated with alkaline phosphatase. The immune reaction was visualized by adding NBT/BCIP in a 1/500 dilution.

The endoGlu-C derived fractions were incubated with one positive serum and one control serum. A strong immunoreaction was revealed with fraction 17. Sequencing of fraction 15 learned that this fraction contained the C-terminal peptide in which the RG motif is dimethylated. Mass analysis of fraction 8 showed that this fraction contained the C-terminal peptide without the modified arginines. Analysis of the fractions located between 8 and 17 indicated that these fractions contain the C-terminal peptide in which the 5 final RG motifs are dimethylated while and the first 4 RG motifs are partially monomethylated. This can be concluded from a mass difference of 14 between the fractions.

The EndoLys-C derived fractions (Fig 1) were incubated separately with 6

positive sera and one control serum. In 5 out of 6 positive sera, a signal significantly higher than the control serum was limited to fraction 15 which was identified both by sequencing and mass analysis as the C-terminal peptide with dimethylarginines. Again, mass analysis indicated that the fractions 8 to 14 correspond to the non-methylated and less methylated forms of the C-terminal peptide.

These results were confirmed by isolating selectively the non-modified and the modified peptide from a preparative endoLys-C digest of baculo SmD1. In Figure 1 it can be seen that fraction 8 with the non-modified SmD1 peptide contained less material than fraction 15 with the dimethylated peptide. It is therefore possible that the exclusive reactivity of the modified peptide was due to different amounts of modified and non-modified peptide being transferred in the dot-blot experiment (Fig.2). To exclude such quantitative variations, the peptides were analysed in a dot spot experiment as described. However, in this experiment equal quantities (based on BCA protein determination) of both peptides, modified and non-modified peptide, were applied. For comparison, the total natural SmD, the total recombinant *E. coli* and baculoviral SmD1 were applied in comparable amounts in the immunodot (Fig 3). Finally peptides were applied in a line immunoassay experiment (Polet *et al.*, Clinical Chemistry, 37, 1991) (Fig.4). Again equal amounts (60 ng) of modified and non-modified SmD1 peptides were applied to a nylon membrane. The amount of peptide bound was visualized by protein colloidal staining (Aurodye, Amersham, Buckinghamshire, UK; Fig. 4). Additionally, 30,15 and 7 ng of recombinant SmD1 from *E.coli*- or baculovirus-infected insect cells as well as a mixture of gel-purified SmD1, SmD2 and SmD3 were applied to the strips. These were then tested with 21 anti-Sm patient sera that were immunoreactive to a mixture of HeLa SmD1, SmD2 and SmD3. Six (29%) of these anti-Sm patient sera gave significant signals with the modified peptide D1, while the non-modified peptide reacted with none of the 21 tested sera. An independent set of anti-Sm Brazilian sera (n=93) showed a comparable rate of reactivity with the modified peptide (4/14 anti-Sm sera; 29%). These experiments substantiate our hypothesis that there are at least 2 epitopes involved in the

immunoreactivity of natural SmD: one epitope is present in the total *E. coli* recombinant SmD molecule while an additional epitope is located in the C-terminus (90-119) of the SmD1 molecule. The presence of dimethylarginines in this peptide is crucial for recognition by patient sera.

5

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
22

References

Andersen,J.,Feeney,R.J. and Zieve,G.W. 1990. Identification and characterization of the small nuclear ribonucleoprotein particle D' core protein. Mol.Cell.Biol. **10**, 4480-4485

Barakat,S.,Briand,J.-P.,Weber,J.-C.,Van Regenmortel,M.H.V. and Muller,S. 1990. Recognition of synthetic peptides of Sm-D autoantigen by lupus sera. Clin.exp.Immunol. **81**, 256-262

De Keyser,F.G., Verbruggen,G., Veys, E.M., Nimmegeers,J., Schatteman,L., Goethals,K., Vandenbossche,M. 1990. "Microgel Diffusionblotting" for sensitive detection of antibodies to extractable nuclear antigens. Clin.Chem. **36**, 337-339

Gheuens,J., MacFarlin,D. 1982. Use of monoclonal anti-idiotypic antibody to P3-X63Ag8 myeloma protein for analysis and purification of B lymphocyte hybridoma products. Eur.J.Immunol. **12**, 701-703

Hoch,S.O. 1989. Application of protein blotting in the study of autoimmune disease. In *Manual of Biological Markers of Disease* B2.4:1-29, Kluwer, Netherlands

Weiner,H.L. 1997. Oral tolerance for the treatment of autoimmune diseases. Annu.Rev.Med. **48**, 341-351

James,J.A.,Mamula,M.J. and Harley,J.B. 1994. Sequential autoantigenic determinants of the small nuclear ribonucleoprotein Sm D shared by human lupus autoantibodies and MRL *lpr/lpr* antibodies. Clin.Exp.Immunol. **98**, 419-426

Lehmeier,T.,Foulaki,K. And Lührman,R. 1990. Evidence for three distinct D proteins, which react differentially with anti-Sm autoantibodies, in the cores of the

major snRNPs U1, U2, U4/U6 and U5. Nucleic Acids Res **18**, 6475-6484

5 Najbauer, J., Johnson, B.A., Young, A.L. and Aswad, D.W. 1993. Peptides with sequences similar to glycine arginine rich motifs in proteins interacting with RNA are efficiently recognized by methyltransferases modifying arginine in numerous proteins. J.Biol.Chem. **268**, 10501-10509

10 Rajpurohit, R., Lee, S.O., Park, J.O., Paik, W.K. and Kim, S. 1994. Enzymatic methylation of recombinant heterogeneous nuclear RNP protein A1. J.Biol.Chem. **269**, 1075-1082

15 Rawal, N., Rajpurohit, R., Lischwe, M.A., Williams, K., R., Paik, W., K., Kim, S. 1995. Structural specificity of substrate for S-Adenosylmethionine:protein arginine N-methyltransferases. Biochem.Biophys.Acta, **1248**, 11-18

Rivkin, E., Vella, M.J. and Lahita, R.G. 1994. A heterogeneous immune response to an Sm-D-like epitope by SLE patients. J.Autoimmun. **7**, 119-132

20 Rokeach, L.A., Haselby, J.A and Hoch, S.O. 1988. Molecular cloning of a cDNA encoding the human Sm-D autoantigen. Proc.Natl.Acad.Sci.USA, **85**, 4832-4836

25 Rokeach, L.A., Haselby, J.A. and Hoch, S.O. 1992a. Overproduction of a human (snRNP)-associated Sm-D autoantigen in *Escherichia coli* and *Saccharomyces cerevisiae*. Gene **118**, 247-253

Rokeach, L.A., Jannatipour, M., Haselby, J.A. and Hoch, S.O. 1992b. Mapping of the immunoreactive domains of a small nuclear ribonucleoprotein-associated Sm-D autoantigen. Clin.Immunol.Immunopath. **35**, 315-324

30 Sabbatini, A., Dolcher, M.P., Marchini, B., Bombardieri, S. And Migliorini, P. 1993a. Mapping of epitopes on the SmD molecule: the use of multiple antigen peptides

to measure autoantibodies in systemic lupus erythematosus. J.Rheumatol. 20,1679-1683

5 Sabbatini,A.,Bombardieri,S. And Migliorini,P. 1993b. Autoantibodies from patients with systemic lupus erythematosus bind a shared sequence of SmD and Epstein-Barr virus-encoded nuclear antigen EBNA I. Eur.J.Immunol. **23**, 1146-1152

10 Van Venrooij,W.J., P. Charles and R.N. Maini 1991. The consensus workshops for the detection of autoantibodies to intracellular antigens in rheumatic diseases. J. Immunol. Methods, **140**, 181-189

15 Wagatsuma,M.,Asami,N.,Miyachi,J.,Uchida,S.,Watanabe,H. and Amann,E. 1993. Antibody recognition of the recombinant human nuclear antigens RNP 70 kD, Sm-A, Sm-B and Sm-D by autoimmune sera. Mol.Immunol. **30**, 1491-1498

Claims

5

- 10

15

10

Figure 1

100

55
56

20

2.

3

an analog of said peptides comprising conservative amino acid substitutions.

3) Peptide and/or chemical structure comprising any of the peptides according to claims 1 or 2, fused to a linker molecule.

4) Circularized peptide that comprises at least one of the peptides according to any of the claims 1 to 3.

5) Peptide comprising and/or consisting of tandem repeats of at least two of any of the peptides of claims 1 to 4.

6) Branched peptide that comprises at least one of the peptides according to any of the claims 1 to 5.

7) Method for producing a peptide according to any of claims 1 to 6, by classical chemical synthesis, wherein methylated arginines are substituted for unmethylated arginine residues during the chemical synthesis.

8) Method for producing a peptide according to any of claims 1 to 6, wherein the primary amino acid sequence is produced by classical chemical synthesis, and wherein the arginine residues that precede glycine residues are subsequently methylated by contacting said peptide with a protein arginine methyltransferase.

9) Method for producing a peptide of any of claims 1 to 6 comprising the following steps:

-transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements such that said peptide or a protein comprising said peptide is expressed and/or secreted,

-culturing said transformed cellular host under conditions allowing expression of said protein or peptide and allowing a partial or optimal methylation of the arginines present in said peptide,

-harvesting said peptide.

10) Method for producing a peptide of any of claims 1 to 6 comprising the following steps:

- transforming an appropriate cellular host with a recombinant vector in which a polynucleic acid is inserted comprising the sequence that codes for said peptide under the control of the appropriate regulatory elements, such that said peptide or a protein comprising said peptide is expressed and/or secreted,
- culturing said transformed cellular host under conditions allowing expression of said protein or said peptide,
- harvesting said protein or said peptide,
- methylating arginine residues of said protein or said peptide by contacting with a protein arginine methyltransferase.

11) Method according to any of claims 9 or 10, wherein said host cell is a bacterial host or yeast or any other eukaryotic host cell which is preferably transformed with a recombinant baculovirus.

12) An antibody raised upon immunization with a peptide according to any of the claims 1 to 6, with said antibody being specifically reactive with the methylated forms of said peptide, and with said antibody being preferably a monoclonal antibody.

13) Anti-idiotypic antibody raised upon immunization with an antibody according to claim 12, with said anti-idiotypic antibody being specifically reactive with the antibody of claim 12, thereby mimicking the methylated form of a peptide according to any of claims 1 to 6, and with said antibody being preferably a monoclonal antibody.

14) An immunotoxin molecule comprising and/or consisting of cell recognition molecule being a peptide of any of claims 1 to 6, or an antibody according to any of the claims 12 or 13, covalently bound to a toxin molecule or active fragment thereof.

15) A peptide according to any of the claims 1 to 6 or an antibody according to any of claims 12 or 13 or an immunotoxin molecule according to claim 14 or a composition thereof for use as a medicament.

5 16) Use of a peptide according to any of claims 1 to 6 or an antibody according to any of claims 12 or 13 or an immunotoxin molecule according to claim 14 or a composition thereof for the preparation of a medicament or of a diagnosticum for auto-immune diseases such as:

- systemic lupus erythematosus,
- 10 -discoid lupus erythematosus,
- scleroderma,
- dermatomyositis,
- rheumatoid arthritis,
- Sjögren's syndrome.

15 or for diseases in which Epstein-Barr can be implicated such as:

- Burkitt's lymphoma,
- nasopharyngeal carcinoma,
- infectious, recurrent or chronic mononucleosis,

20 17) Use of a polypeptide according to claim 1 to 6 or a composition thereof for the preparation of a medicament to treat auto-immune diseases by increasing the size of antigen-immune complexes, thereby improving the clearance of the formed immune complexes.

25 18) Use of a polypeptide according to claim 1 to 6 or a composition thereof for the preparation of a medicament for oral administration to treat auto-immune diseases by inducing a state of systemic hyporesponsiveness to the said polypeptide ('Oral tolerance').

30 19) A diagnostic kit for use in detecting auto-immune diseases such as:

- systemic lupus erythematosus,

- discoid lupus erythematosus,
- scleroderma,
- dermatomyositis,
- rheumatoid arthritis,
- Sjögren's syndrome,

or for detecting diseases in which Epstein-Barr can be implicated such as:

- Burkitt's lymphoma,
- nasopharyngeal carcinoma,
- Hodgkin's disease,
- infectious, recurrent or chronic mononucleosis,

said kit comprising at least one peptide according to any of claims 1 to 6, or an antibody according to claims 12 or 13, with said peptide or antibody being possibly bound to a solid support.

20) A diagnostic kit according to claim 19, said kit comprising a range of peptides according to any of claims 1 to 6 or of antibodies according to claims 12 or 13, possibly in combination with native methylated SmD1 or SmD3 and recombinant unmethylated SmD1 or SmD3, wherein said peptides are attached to specific locations on a solid substrate.

21) A diagnostic kit according to claim 19 or 20, wherein said solid support is a membrane strip and said polypeptides are coupled to the membrane in the form of parallel lines.

- natural SmD (1,2 or 3) or in vitro dimethylated SmD (1,2 or 3)
- unmethylated SmD expressed in E.coli (1,2 or 3)
- peptide of any of claims 1 to 6

22) A diagnostic kit according to any of claims 19 to 21 wherein certain peptides are not attached to a solid support but are provided in the binding solution to be used as competitors and/or to block other antibodies that are present in sera from patients with autoimmune disease other than SLE, thereby decreasing or

eliminating possible cross-reaction and/or aspecific binding.

the following examples are given for illustration purposes only and are not intended to limit the scope of the invention.




Figure 1




Fig. 2

3/4




Fig. 3




Fig. 4

D E C L A R A T I O N

As below named inventors, we hereby declare that:

Our residence, post office address and citizenship are as stated below next to our names.

The below named inventors are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHYLATED, S_mD HOMOLOGOUS PEPTIDES, REACTIVE WITH THE ANTIBODIES FROM SERA OF LIVING BEINGS AFFECTED WITH SYSTEMIC LUPUS ERYTHEMATOSUS**, the specification of which was filed as PCT International Application No. PCT/EP98/05518 on August 31, 1998 and was not amended under PCT Article 19.

We hereby state that we have reviewed and understand the contents of the above identified specification, including the claims.

We acknowledge the duty to disclose to the Patent and Trademark Office all information known to us to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

We hereby claim foreign priority benefits under Title 35, United States Code, § 119 (a)-(d) of any foreign application(s) for patent listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

PRIOR FOREIGN APPLICATION(S)

Priority Claimed

<u>97870127.4</u>	<u>Europe</u>	<u>29 August 1997</u>	<u>Yes</u>
(Number)	(Country)	(Date Filed)	

We hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, we acknowledge the duty to disclose all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56, which become available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>PCT/EP98/05518</u>	<u>31 August 1998</u>
(International Application No.)	(International Filing Date)


We hereby direct that all correspondence and telephone calls be addressed to:

Patricia A. Kammerer
Arnold, White & Durkee
P. O. Box 4433
Houston, Texas 77210-4433
(713) 787-1438

attorneys for the prospective assignee of this application.


WE HEREBY DECLARE THAT ALL STATEMENTS MADE OF OUR OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH, UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.

100


Inventor's Full Name	<u>LYDIE</u> <u>MEHEUS</u>	
Inventor's Signature		
Date: <u>23 / 4 / 99</u>	Country of Citizenship: <u>Belgium</u>	
Residence Address	120 Gontrodeheirweg B-9820 <u>Merelbeke</u> <u>Belgium</u> <u>BEX</u>	
Post Office Address, if different from above	same as above	

Inventor's Full Name	<u>REINHARD</u> <u>GEORG</u> <u>LÜHRMANN</u>
Inventor's Signature	
Date:	Country of Citizenship: <u>Germany</u>
Residence Address	
Post Office Address, if different from above	same as above

200

Inventor's Full Name	<u>ANN</u> <u>UNION</u>
Inventor's Signature	
Date: <u>23 / 4 / 99</u>	Country of Citizenship: <u>Belgium</u>
Residence Address	15 Savooien B-9880 <u>Aalter</u> <u>Belgium</u> <u>BEX</u>
Post Office Address, if different from above	same as above

300

Inventor's Full Name	<u>JOSEPH</u> <u>RAYMACKERS</u>
Inventor's Signature	
Date: <u>23 / 4 / 99</u>	Country of Citizenship: <u>Belgium</u>
Residence Address	10 Zuiderbiesten B-9810 <u>Eke</u> <u>Belgium</u> <u>BEX</u>
Post Office Address, if different from above	same as above

D E C L A R A T I O N

As below named inventors, we hereby declare that:

Our residence, post office address and citizenship are as stated below next to our names.

The below named inventors are the original, first and joint inventors of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHYLATED, SmD HOMOLOGOUS PEPTIDES, REACTIVE WITH THE ANTIBODIES FROM SERA OF LIVING BEINGS AFFECTED WITH SYSTEMIC LUPUS ERYTHEMATOSUS**, the specification of which was filed as PCT International Application No. PCT/EP98/05518 on August 31, 1998 and was not amended under PCT Article 19.

We hereby state that we have reviewed and understand the contents of the above identified specification, including the claims.

We acknowledge the duty to disclose to the Patent and Trademark Office all information known to us to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

We hereby claim foreign priority benefits under Title 35, United States Code, § 119 (a)-(d) of any foreign application(s) for patent listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed.

<u>PRIOR FOREIGN APPLICATION(S)</u>			<u>Priority Claimed</u>
<u>97870127.4</u>	<u>Europe</u>	<u>29 August 1997</u>	<u>Yes</u>
(Number)	(Country)	(Date Filed)	

We hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT International application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code, § 112, we acknowledge the duty to disclose all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56, which become available between the filing date of the prior application and the national or PCT international filing date of this application.

<u>PCT/EP98/05518</u>	<u>31 August 1998</u>
(International Application No.)	(International Filing Date)


We hereby direct that all correspondence and telephone calls be addressed to:

Patricia A. Kammerer
Arnold, White & Durkee
P. O. Box 4433
Houston, Texas 77210-4433
(713) 787-1438

attorneys for the prospective assignee of this application.


WE HEREBY DECLARE THAT ALL STATEMENTS MADE OF OUR OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH, UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.

10


Inventor's Full Name	<u>LYDIE</u> <u>MEHEUS</u>	
Inventor's Signature		
Date: <u>23 / 4 / 99</u>	Country of Citizenship:	<u>Belgium</u>
Residence Address	120 Gontrodeheirweg B-9820 <u>Merelbeke</u> Belgium BEX	
Post Office Address, if different from above	same as above	

Inventor's Full Name	<u>REINHARD</u> <u>GEORG</u> <u>LÜHRMANN</u>
Inventor's Signature	
Date:	Country of Citizenship: <u>Germany</u>
Residence Address	
Post Office Address, if different from above	same as above

20

Inventor's Full Name	<u>ANN</u> <u>UNION</u>
Inventor's Signature	
Date: <u>23 / 4 / 99</u>	Country of Citizenship: <u>Belgium</u>
Residence Address	15 Savooien B-9880 <u>Aalter</u> Belgium BEX
Post Office Address, if different from above	same as above

30

Inventor's Full Name	<u>JOSEPH</u> <u>RAYMACKERS</u>
Inventor's Signature	
Date: <u>23 / 4 / 99</u>	Country of Citizenship: <u>Belgium</u>
Residence Address	10 Zuiderbiesten B-9810 <u>Eke</u> Belgium BEX
Post Office Address, if different from above	same as above